

Baryon Acoustic Oscillations in the Ly α forest of BOSS DR11 quasars

Timothée Delubac^{1,12}, Julian E. Bautista², Nicolás G. Busca^{2,24,25}, James Rich¹, David Kirkby³, Stephen Bailey⁴, Andreu Font-Ribera⁴, Anže Slosar⁵, Khee-Gan Lee⁶, Matthew M. Pieri⁷, Jean-Christophe Hamilton², Éric Aubourg², Michael Blomqvist³, Jo Bovy⁸, J. Brinkmann⁹, William Carithers⁴, Kyle S. Dawson¹⁰, Daniel J. Eisenstein¹¹, Jean-Paul Kneib^{12,13}, J.-M. Le Goff¹, Daniel Margala³, Jordi Miralda-Escudé^{14,15}, Adam D. Myers¹⁶, Robert C. Nichol⁷, Pasquier Noterdaeme¹⁷, Ross O’Connell¹⁸, Matthew D. Olmstead¹⁰, Nathalie Palanque-Delabrouille¹, Isabelle Pâris¹⁷, Patrick Petitjean¹⁷, Nicholas P. Ross^{4,19}, Graziano Rossi^{1,26}, David J. Schlegel⁴, Donald P. Schneider^{20,21}, David H. Weinberg²², Christophe Yèche¹, Donald G. York²³

¹ CEA, Centre de Saclay, IRFU, F-91191 Gif-sur-Yvette, France

² APC, Université Paris Diderot-Paris 7, CNRS/IN2P3, CEA, Observatoire de Paris, 10, rue A. Domon & L. Duquet, Paris, France

³ Department of Physics and Astronomy, University of California, Irvine, CA 92697, USA

⁴ Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA

⁵ Brookhaven National Laboratory, 2 Center Road, Upton, NY 11973, USA

⁶ Max-Planck-Institut für Astronomie, Königstuhl 17, D69117 Heidelberg, Germany

⁷ Institute of Cosmology and Gravitation, Dennis Sciama Building, University of Portsmouth, Portsmouth, PO1 3FX, UK

⁸ Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA

⁹ Apache Point Observatory, P.O. Box 59, Sunspot, NM 88349, USA

¹⁰ Department of Physics and Astronomy, University of Utah, 115 S 1400 E, Salt Lake City, UT 84112, USA

¹¹ Harvard-Smithsonian Center for Astrophysics, Harvard University, 60 Garden St., Cambridge MA 02138, USA

¹² Laboratoire d’Astrophysique, Ecole polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland

¹³ Aix Marseille Université, CNRS, LAM (Laboratoire d’Astrophysique de Marseille) UMR 7326, F-13388, Marseille, France

¹⁴ Institució Catalana de Recerca i Estudis Avançats, Barcelona, Catalonia

¹⁵ Institut de Ciències del Cosmos, Universitat de Barcelona (UB-IEEC), Catalonia

¹⁶ Department of Physics and Astronomy, University of Wyoming, Laramie, WY 82071, USA

¹⁷ Université Paris 6 et CNRS, Institut d’Astrophysique de Paris, 98bis blvd. Ara go, 75014 Paris, France

¹⁸ Bruce and Astrid McWilliams Center for Cosmology, Carnegie Mellon University, Pittsburgh, PA 15213, USA

¹⁹ Dept. of Physics, Drexel University, 3141 Chestnut Street, Philadelphia, PA 19104, USA

²⁰ Department of Astronomy and Astrophysics, The Pennsylvania State University, University Park, PA 16802

²¹ Institute for Gravitation and the Cosmos, The Pennsylvania State University, University Park, PA 16802

²² Department of Astronomy, Ohio State University, 140 West 18th Avenue, Columbus, OH 43210, USA

²³ Department of Astronomy and Astrophysics and the Enrico Fermi Institute, The University of Chicago, 5640 South Ellis Avenue, Chicago, Illinois, 60615, USA

²⁴ Observatório Nacional, Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil

²⁵ Laboratório Interinstitucional de e-Astronomia, - LIneA, Rua Gal. José Cristino 77, Rio de Janeiro, RJ - 20921-400, Brazil

²⁶ Department of Astronomy and Space Science, Sejong University, Seoul, 143-747, Korea

Received April 1, 2012; accepted April 1, 2014

ABSTRACT

We report a detection of the baryon acoustic oscillation (BAO) feature in the flux-correlation function of the Ly α forest of high-redshift quasars with a statistical significance of five standard deviations. The study uses 137,562 quasars in the redshift range $2.1 \leq z \leq 3.5$ from the Data Release 11 (DR11) of the Baryon Oscillation Spectroscopic Survey (BOSS) of SDSS-III. This sample contains three times the number of quasars used in previous studies. The measured position of the BAO peak determines the angular distance, $D_A(z = 2.34)$ and expansion rate, $H(z = 2.34)$, both on a scale set by the sound horizon at the drag epoch, r_d . We find $D_A/r_d = 11.28 \pm 0.65(1\sigma)_{-1.2}^{+2.8}(2\sigma)$ and $D_H/r_d = 9.18 \pm 0.28(1\sigma) \pm 0.6(2\sigma)$ where $D_H = c/H$. The optimal combination, $\sim D_H^{0.7} D_A^{0.3}/r_d$ is determined with a precision of $\sim 2\%$. For the value $r_d = 147.4$ Mpc, consistent with the CMB power spectrum measured by Planck, we find $D_A(z = 2.34) = 1662 \pm 96(1\sigma)$ Mpc and $H(z = 2.34) = 222 \pm 7(1\sigma)$ km s⁻¹Mpc⁻¹. Tests with mock catalogs and variations of our analysis procedure have revealed no systematic uncertainties comparable to our statistical errors. Our results agree with the previously reported BAO measurement at the same redshift using the quasar-Ly α forest cross-correlation. The auto-correlation and cross-correlation approaches are complementary because of the quite different impact of redshift-space distortion on the two measurements. The combined constraints from the two correlation functions imply values of D_A/r_d and D_H/r_d that are, respectively, 7% low and 7% high compared to the predictions of a flat Λ CDM cosmological model with the best-fit Planck parameters. With our estimated statistical errors, the significance of this discrepancy is $\approx 2.5\sigma$.

Key words. cosmology, Ly α forest, large scale structure, dark energy

1. Introduction

Observation of the peak in the matter correlation function due to Baryon Acoustic Oscillations (BAO) in the pre-recombination epoch is now an established tool to constrain cosmological models. The BAO peak at a redshift z appears at an angular separation $\Delta\theta = r_d/[(1+z)D_A(z)]$ and at a redshift separation $\Delta z = r_d/D_H(z)$, where D_A and $D_H = c/H$ are the angular and Hubble distances and r_d is the sound horizon at the drag epoch¹. Measurement of the peak position at any redshift thus constrains the combinations of cosmological parameters that determine D_H/r_d and D_A/r_d .

The BAO peak has been observed primarily in the galaxy-galaxy correlation function obtained in redshift surveys. The small statistical significance of the first studies gave only constraints on D_V/r_d where D_V is the combination $D_V = [(1+z)D_A]^{2/3}[zD_H]^{1/3}$ which determines the peak position for the galaxy correlation function when averaged over directions with respect to the line of sight. The first measurements were at $z \sim 0.3$ by the SDSS (Eisenstein et al., 2005) and 2dFGRS (Cole et al., 2005) with results from the combined data set presented by Percival et al. (2010). A refined analysis using reconstruction (Eisenstein et al., 2007; Padmanabhan et al., 2009) to sharpen the precision D_V/r_d was presented by Padmanabhan et al. (2012) and Mehta et al. (2012).

Other measurements of D_V/r_d were made at $z \sim 0.1$ by the 6dFGRS (Beutler et al., 2011), at $(0.4 < z < 0.8)$ by WiggleZ (Blake et al., 2011a), and, using galaxy clusters, at $z \sim 0.3$ by Veropalumbo et al. (2013). The Baryon Oscillation Spectroscopic Survey (BOSS; Dawson et al. 2013) of SDSS-III (Eisenstein et al., 2011) has presented measurements of D_V/r_d at $z \sim 0.57$ and $z \sim 0.32$ (Anderson et al., 2012). A measurement at $z \sim 0.54$ of D_A/r_d using BOSS photometric data was made by Seo et al. (2012).

The first combined constraints on D_H/r_d and D_A/r_d were obtained using the $z \sim 0.3$ SDSS data by Chuang & Wang (2012) and Xu et al. (2012). Recently, BOSS has provided precise constraints on D_H/r_d and D_A/r_d at $z = 0.57$ (Anderson et al., 2013; Kazin et al., 2013).

At higher redshifts, the BAO feature can be observed using absorption in the Ly α forest to trace mass, as suggested by McDonald (2003), White (2003) and McDonald & Eisenstein (2007). After the observation of the predicted large scale correlations in early BOSS data by Slosar et al. (2011), a BAO peak in the Ly α forest correlation function was measured by BOSS in the SDSS data release DR9 (Busca et al., 2013; Slosar et al., 2013; Kirkby et al., 2013). The peak in the quasar-Ly α forest cross correlation function was detected in the larger data sets of DR11 (Font-Ribera et al., 2013) and DR10 (O’Connell et al., in preparation). The DR10 data are now public (Ahn et al., 2013), and the DR11 data will be made public simultaneously with the final SDSS-III data release (DR12) in late 2014.

This paper presents a new measurement of the Ly α forest auto-correlation function and uses it to study BAO at $z = 2.34$. It is based on the methods used by Busca et al. (2013) but introduces several improvements in the analysis. First, and most im-

portant, is a tripling of the number of quasars by using the DR11 catalog of 158,401 quasars in the redshift range $2.1 \leq z_q \leq 3.5$. Second, to further increase the statistical power we use a slightly expanded forest range as well as quasars that have damped Ly α troughs in the forest. Finally, the Busca et al. (2013) analysis was based on a decomposition of the correlation function into monopole and quadrupole components. Here, we fit the full correlation $\xi(r_\perp, r_\parallel)$ as a function of separations perpendicular, r_\perp , and parallel, r_\parallel , to the line of sight. This more complete treatment is made possible by a more careful determination of the covariance matrix compared to that of Busca et al. (2013).

Our analysis uses a fiducial cosmological model in two places. First, flux pixel pairs separated in angle and wavelength are assigned a co-moving separation (in h^{-1} Mpc) using the $D_A(z)$ and $D_H(z)$ calculated with the adopted parameters. Second, to determine the observed peak position, we compare our measured correlation function with a correlation function generated using CAMB (Lewis et al., 2000) as described in Kirkby et al. (2013). We adopt the same (flat) Λ CDM model used in Busca et al. (2013), Slosar et al. (2013) and Font-Ribera et al. (2013); with the parameters given in table 1. The fiducial model has values of D_A/r_d and D_H/r_d at $z = 2.34$ that differ by of order 1% from the values given by the models favored by CMB data (Ade et al., 2013; Calabrese et al., 2013) given in the second and third columns of table 1.

This paper is organized as follows. Section 2 describes the DR11 data used in this analysis. Section 3 gives a brief description of the mock spectra used to test the analysis procedure, with a more detailed description being found in Bautista et al. (in preparation). Section 4 presents our method of estimating the correlation function $\xi(r_\perp, r_\parallel)$ and its associated covariance matrix. In section 5 we fit the data to derive the BAO peak position parameters, $D_A(z = 2.34)/r_d$ and $D_H(z = 2.34)/r_d$. Section 6 investigates possible systematic errors in the measurement. In section 7 we compare our measured peak position to that measured by the Quasar-Ly α -forest cross-correlation (Font-Ribera et al., 2013) and study Λ CDM models that are consistent with these results. Section 8 concludes.

2. The BOSS quasar sample and data reduction

The BOSS project (Dawson et al., 2013) of SDSS-III (Eisenstein et al., 2011) was designed to obtain the spectra of over $\sim 1.6 \times 10^6$ luminous galaxies and $\sim 150,000$ quasars. The project uses upgraded versions of the SDSS spectrographs (Smee et al., 2012) mounted on the Sloan 2.5-meter telescope (Gunn et al., 2006) at Apache Point, New Mexico.

The quasar spectroscopy targets are selected from photometric data via a combination of algorithms (Richards et al., 2009; Yeche et al., 2009; Kirkpatrick et al., 2011; Bovy et al., 2011; Palanque-Delabrouille et al., 2011) as summarized in Ross et al. (2012). The algorithms use SDSS ugriz fluxes (Fukugita et al., 1996; York et al., 2000) and, for SDSS Stripe 82, photometric variability. Using the techniques of Bovy et al. (2012), we also use any available data from non-optical surveys: the GALEX survey (Martin et al., 2005) in the UV; the UKIDSS survey (Lawrence et al., 2007) in the NIR, and the FIRST survey (Becker et al., 1995) in the radio.

In this paper we use the data from the DR11 data release of SDSS-III, whose footprint is shown in figure 1. These data cover 8377 deg^2 of the ultimate BOSS 10^4 deg^2 footprint.

The data were reduced with the SDSS-III pipeline as described in Bolton et al. (2012). Typically, four exposures of 15 minutes were co-added in pixels of wavelength width $\Delta \log_{10} \lambda =$

¹ We follow the convention of Anderson et al. (2013), $r_d = r_s(z_d)$, where r_s is the sound horizon and z_d is the drag redshift (baryon decoupling from photons), to be distinguished from z_* (the redshift corresponding to unity optical depth for CMB photons). Earlier publications on BAO generally denoted r_d simply as r_s . For models with cold dark matter, baryons and three light neutrino species, r_d can be evaluated with equation (55) of Anderson et al. (2013), which agrees with the CAMB-derived value to better than 0.1 per cent.




Fig. 1. Hammer-Aitoff projection of the BOSS DR11 footprint (dec. vs. r.a.) used here. The light areas show the DR9 sub-region available for the earlier studies of Busca et al. (2013) and Slosar et al. (2013). The red-dashed line shows the location of the galactic plane.

Table 1. Parameters of the fiducial flat Λ CDM cosmological model used for this analysis, the flat Λ CDM model derived from Planck and low- ℓ WMAP polarisation data, ‘Planck + WP’ (Ade et al., 2013), and a flat Λ CDM model derived from the WMAP, ACT and SPT data (Calabrese et al., 2013). The models are defined by the cold dark matter, baryon and massive neutrinos densities, the Hubble constant and the number of light neutrino species. The sound horizon at the drag epoch, r_d is calculated using CAMB (which can be approximated with equation (55) of Anderson et al. (2013) to a precision of 0.1%).

	fiducial	Planck + WP	WMAP9 +ACT+SPT
$\Omega_M h^2$	0.1323	0.14305	0.1347
$= \Omega_C h^2$	0.1090	0.12038	0.1122
$+ \Omega_B h^2$	0.0227	0.022032	0.02252
$+ \Omega_\nu h^2$	0.0006	0.0006	0
h	0.7	0.6704	0.714
N_ν	3	3	3
Ω_M	0.27	0.3183	0.265
r_d (Mpc)	149.7	147.4	149.1
	(104.80 h^{-1})	(98.79 h^{-1})	(106.4 h^{-1})
$D_A(2.34)/r_d$	11.59	11.76	11.47
$D_H(2.34)/r_d$	8.708	8.570	8.648

10^{-4} ($c\Delta\lambda/\lambda \sim 69 \text{ km s}^{-1}$). The pipeline provides flux calibrated spectra, object classifications (galaxy, quasar, star), and redshift estimates for all targets.

The spectra of all quasar targets were visually inspected (Pâris et al., 2012, 2013) to correct for misidentifications or inaccurate redshift determinations and to flag broad absorption lines (BALs). Damped Ly α troughs were visually flagged, but also identified and characterized automatically (Noterdaeme et al., 2012). The visual inspection of DR11 confirmed 158,401

quasars with $2.1 \leq z_q \leq 3.5$. In order to simplify the analysis of the Ly α forest, we discarded quasars with visually identified BALs, leaving 140,579 quasars. A further cut requiring a minimum number of unmasked forest pixels (50 ‘analysis pixels’; see below) yielded a sample of 137,562 quasars.

For the measurement of the flux transmission, we use the rest-frame wavelength interval

$$104.0 < \lambda_{\text{rf}} < 120.0 \text{ nm}, \quad (1)$$

slightly wider than in Busca et al. (2013). This range is bracketed by the Ly β and Ly α emission lines at 102.5 and 121.6 nm, and was chosen as the maximum range that avoids the large pixel variances on the slopes of the two lines due to quasar-to-quasar diversity of line-emission strength. The absorber redshift, $z = \lambda/\lambda_{\text{Ly}\alpha} - 1$, is required to be in the range $1.96 < z < 3.44$. The lower limit is set by the requirement that the observed wavelength be greater than 360 nm, below which the system throughput is less than 10% its peak value. The upper limit is produced by the maximum quasar redshift of 3.5, beyond which the BOSS surface density of quasars is not sufficient to be useful for this study. The weighted distribution of redshifts of absorber pairs near the BAO peak position is shown in figure 2 (top panel); it has a mean of $\langle z \rangle = 2.34$.

Forests with identified DLAs are given a special treatment. All pixels where the absorption due to the DLA is higher than 20% are not used. Otherwise, the absorption in the wings is corrected using a Voigt profile following the procedure of Noterdaeme et al. (2012). The metal lines due to absorption at the DLA redshift are masked. The lines to be masked were those identified in a stack of spectra shifted to the redshift of the detected DLA. The width of the mask was 0.2 nm or 0.3 nm (depending on the line strength) or 4.1 nm for Ly β . We also mask the ± 3 nm region corresponding to Ly α if the DLA finder erroneously interpreted Ly β absorption as Ly α absorption.

For the determination of the correlation function, we use ‘analysis pixels’ that are the flux average over three adja-




Fig. 2. Top: distribution of redshifts of pairs of pixels contributing to ξ in the region $80 < r < 120 h^{-1}\text{Mpc}$. Bottom: distribution of all pixel redshifts

cent pipeline pixels. Throughout the rest of this paper, “pixel” refers to analysis pixels unless otherwise stated. The width of these pixels is 207 km s^{-1} , i.e. an observed-wavelength width $\sim 0.27 \text{ nm}$ or $\sim 2 h^{-1}\text{Mpc}$. The total sample of 137,562 quasars thus provides $\sim 2.4 \times 10^7$ measurements of Ly α absorption over a total volume of $\sim 50 h^{-3}\text{Gpc}^3$.

3. Mock quasar spectra

In addition to the BOSS spectra, we analyzed 100 sets of mock spectra. This exercise was undertaken to search for possible systematic errors in the recovered BAO peak position and to verify that uncertainties in the peak position are correctly estimated. The spectra were generated using the methods of Font-Ribera et al. (2012a). A detailed description of the production and resulting characteristics of the mock spectra is given in Bautista et al. (in preparation).

For each set of spectra, the background quasars were assigned the angular positions and redshifts of the DR11 quasars. The foreground absorption field in redshift space was generated according to a cosmology similar to the fiducial cosmol-




Fig. 3. The measured correlation function averaged over three angular regions: $\mu > 0.8$ (top), $0.8 > \mu > 0.5$ (middle), and $0.5 > \mu > 0.0$ (bottom), where μ is the central value of $r_{\parallel}/\sqrt{r_{\parallel}^2 + r_{\perp}^2}$ in each $(r_{\parallel}, r_{\perp})$ bin. The gray lines show individual sets of mocks and the solid blue line represents the mean of the 100 mock sets. The dashed blue lines are the 1σ variations of the mocks. The red points show the data.

ogy of Table 1.² The unabsorbed spectra (continua) of the quasars were generated using the Principal Component Analysis (PCA) eigenspectra of Suzuki et al. (2005), with amplitudes for each eigenspectrum randomly drawn from Gaussian distri-

² The model has the same $\Omega_M h^2$ but $\Omega_b h^2 = 0$. This change has a negligible impact on the generated power spectrum and changes the BAO peak position by only 0.1%.

butions with sigma equal to the corresponding eigenvalues as published in Table 1 of Suzuki (2006). Finally, the spectra were modified to include the effects of the BOSS spectrograph point spread function (PSF), readout noise, photon noise, and flux systematic errors.

Our mock production pipeline admits the option of adding DLAs to the spectra according to the procedure described in Font-Ribera et al. (2012b). However, since identified DLAs are masked in the analysis of real data, we did not simulate them into the mocks. Of course low column density and Lyman limit systems are not efficiently identified and masked in the data, so such systems are present in the data but not in the mocks.

Absorption by metals was added to a separate group of ten mocks according to the procedure described in Bautista et al. (in preparation). The quantity of each metal to be added was determined by a modified Ly α stacking procedure from Pieri et al. (2010) and Pieri et al. (2013). As discussed in Section 6, the metals have an effect on the recovered correlation function only at small transverse separations, $r_{\perp} < 10 h^{-1}\text{Mpc}$, and have no significant effect on the measured position of the BAO peak.

A total of 100 independent metal-free realizations of the BOSS data were produced and analyzed with the same procedures as those for the real data. Figure 3 shows the correlation function of the mocks and the data as measured by the techniques described in the next section. The mocks reproduce well the general features of the observed correlation function. We therefore use them in section 5.2 to search for biases in the analysis procedure that would influence the position of the BAO peak.

4. Measurement of the correlation function

In this section we describe the measurement of the correlation function of the transmitted flux fraction:

$$\delta_q(\lambda) = \frac{f_q(\lambda)}{C_q(\lambda)\bar{F}(z)} - 1. \quad (2)$$

Here, $f_q(\lambda)$ is the observed flux density for quasar q at observed wavelength λ , $C_q(\lambda)$ is the unabsorbed flux density (the so-called ‘‘continuum’’) and $\bar{F}(z)$ is the mean transmitted fraction at the absorber redshift, $z(\lambda) = \lambda/\lambda_{\text{Ly}\alpha} - 1$. Figure 4 shows a spectrum with its $C_q(\lambda)$ (blue line) and $C_q\bar{F}$ (red line) estimated by the methods of section 4.1.

For the estimator of the correlation function, we use a simple weighted sum of products of the deltas:

$$\hat{\xi}_A = \frac{\sum_{ij \in A} w_{ij} \delta_i \delta_j}{\sum_{ij \in A} w_{ij}}, \quad (3)$$

where the w_{ij} are weights (section 4.2) and each i or j indexes a measurement on a quasar q at wavelength λ . The sum over (i, j) is understood to run over all pairs of pixels within a bin A in the space of pixel separations, $\mathbf{r}_i - \mathbf{r}_j$. The bins A are defined by a range of width $4 h^{-1}\text{Mpc}$ of the components perpendicular and parallel to the line of sight, r_{\perp} and r_{\parallel} . We use 50 bins in each component, spanning the range from 0 to $200 h^{-1}\text{Mpc}$; the total number of bins used for evaluating the correlation function is therefore 2500. Separations in observational pixel coordinates (ra,dec,z) are transformed to $(r_{\perp}, r_{\parallel})$ in units of $h^{-1}\text{Mpc}$ by using a ΛCDM fiducial cosmology described in table 1.

In the sum (3), we exclude pairs of pixels from the same quasar to avoid the correlated errors in δ_i and δ_j arising from the estimate of $C_q(\lambda)$ for the spectrum of the quasar. The weights in eq. (3) are set to zero for pixels flagged by the pipeline as

having problems due, e.g., to sky emission lines or cosmic rays. We also do not use pairs of pixels that have nearly the same wavelength ($r_{\parallel} < 4 h^{-1}\text{Mpc}$) and that were taken on the same focal-plane plate. The reason for this decision is that these pairs have $\sim 20\%$ greater correlation than expected from our linear cosmological model fit using data with $r_{\parallel} > 4 h^{-1}\text{Mpc}$. This result is most likely to be due to spurious correlations introduced by the pipeline, e.g., from sky subtraction for flux calibration operations.

4.1. Continuum fits

We have used three methods to estimate $C_q\bar{F}$ used in eq. (2). The first two assume that $C_q\bar{F}$ is, to first approximation, the product of two factors: a scaled universal quasar spectrum that is a function of rest-frame wavelength, $\lambda_{\text{rf}} = \lambda/(1+z_q)$ (for quasar redshift z_q), and a mean transmission fraction that slowly varies with absorber redshift. The universal spectrum is found by stacking the appropriately normalized spectra of quasars in our sample, thus averaging the fluctuating Ly α absorption. The continuum for individual quasars is then derived from the universal spectrum by normalizing it to the quasar’s mean forest flux and then modifying its slope to account for spectral-index diversity and/or photo-spectroscopic miscalibration.

Our simplest continuum estimator, C1, is the ‘‘method 1’’ of Busca et al. (2013). It estimates directly the product $C_q\bar{F}$ in equation 2, by modeling each spectrum as

$$C_q\bar{F} = a_q \left(\frac{\lambda}{\langle \lambda \rangle} \right)^{b_q} \bar{f}(\lambda_{\text{rf}}, z), \quad (4)$$

where a_q is a normalization, b_q a ‘‘deformation parameter’’, $\langle \lambda \rangle$ the mean wavelength in the forest for the quasar q , and $\bar{f}(\lambda_{\text{rf}}, z)$ is the mean normalized flux obtained by stacking spectra in bins of width $\Delta z = 0.1$.

As noted in Busca et al. (2013), the mean value of $\delta_q(\lambda)$ (averaged over all measurements in narrow bins in λ) has peaks at the position of the Balmer series of amplitude ~ 0.02 . These artifacts are due to imperfect use of spectroscopic standards containing those lines. These artifacts are removed on average by subtracting the mean δ from each measurement: $\delta_q(\lambda) \rightarrow \delta_q(\lambda) - \langle \delta(\lambda) \rangle$.

The C1 continuum estimator would be close to optimal if the distribution of δ about zero was Gaussian. Since the true distribution is quite asymmetric, we developed a slightly more sophisticated continuum estimator, ‘‘method 2’’ of Busca et al. (2013), denoted here as C2. We adopt it as the standard estimator for this work. The continuum for each quasar is assumed to be of the form

$$C_q(\lambda) = [a_q + b_q \log(\lambda)] \bar{C}(\lambda_{\text{rf}}), \quad (5)$$

where $\bar{C}(\lambda_{\text{rf}})$ is the mean continuum determined by stacking spectra. The parameters a_q and b_q are fitted to match the quasar’s distribution of transmitted flux to an assumed probability distribution derived from the log-normal model used to generate mock data.

The C2 continuum is then multiplied by the mean transmitted flux fraction $\bar{F}(z)$ which we determine by requiring that the mean of the delta field vanish for all redshifts. This last step has the effect of removing the average of the Balmer artifacts.

The third continuum estimate method, C3, is a modified version of the MF-PCA technique described in Lee et al. (2012). This method has been used to provide continua for the publicly




Fig. 4. An example of a BOSS quasar spectrum of redshift 3.239. The red and blue lines cover the forest region used here, $104.0 < \lambda_{\text{rf}} < 120.0$. This region is sandwiched between the quasar’s Ly β and Ly α emission lines respectively at 435 and 515 nm. The blue (green) line is the C2 (C3) model of the continuum, $C_q(\lambda)$, and the red line is the C1 model of the product of the continuum and the mean absorption, $C_q(\lambda)\bar{F}(z)$. (See text.)

available DR9 spectra (Lee et al., 2013). Unlike the other two methods, it does not assume a universal spectral form. Instead, for each spectrum, it fits a variable amplitude PCA template to the part redward of the Ly α wavelength. The predicted spectrum in the forest region is then renormalized so that the mean forest flux matches the mean forest flux at the corresponding redshift.

All three methods use data in the forest region to determine the continuum and therefore necessarily introduce distortions in the flux transmission field and its correlation function (Slosar et al., 2011). Fortunately, these distortions are not expected to shift the BAO peak position, and this expectation is confirmed in the mock spectra.

4.2. Weights

We choose the weights w_{ij} so as to approximately minimize the relative error on $\hat{\xi}_A$ estimated with equation (3). The weights should obviously favor low-noise pixels and take into account the redshift dependence of the pixel correlations, $\xi_{ij}(z) \propto (1 + z_i)^{\gamma/2}(1 + z_j)^{\gamma/2}$, with $\gamma \sim 3.8$ (McDonald et al., 2006). Following Busca et al. (2013), we use

$$w_{ij} \propto \frac{(1 + z_i)^{\gamma/2}(1 + z_j)^{\gamma/2}}{\xi_{ii}^2 \xi_{jj}^2}, \quad (6)$$

where ξ_{ii} is assumed to have noise and LSS contributions:

$$\xi_{ii}^2 = \frac{\sigma_{\text{pipeline},i}^2}{\eta(z_i)} + \sigma_{\text{LSS}}^2(z_i) \quad \text{and} \quad z_i = \lambda_i / \lambda_{\text{Ly}\alpha} - 1. \quad (7)$$

Here $\sigma_{\text{pipeline},i}^2$ is the pipeline estimate of the noise-variance of pixel i multiplied by $(C_i \bar{F}_i)^2$ and η is a factor that corrects for a possible inaccurate estimate of the variance by the pipeline. The two functions $\eta(z)$ and $\sigma_{\text{LSS}}^2(z)$ are determined by measuring the variance of δ_i in bins of $\sigma_{\text{pipeline},i}^2$ and redshift.




Fig. 5. The measured correlation functions (continuum C2) in three angular regions: $\mu > 0.8$ (top), $0.8 > \mu > 0.5$ (middle), and $0.5 > \mu > 0$. (bottom), where μ is the central value of $r_{\parallel} / \sqrt{r_{\parallel}^2 + r_{\perp}^2}$ in each $(r_{\parallel}, r_{\perp})$ bin. The curves show the results of fits as described in section 5. The full curve is best fit and the dashed curve is best fit when the parameters α_{\perp} and α_{\parallel} (equation 11) are both set to unity. The irregularities in the fits are due to the use of $(r_{\parallel}, r_{\perp})$ bins rather than (r, μ) bins.

4.3. $\xi(r_{\perp}, r_{\parallel})$ and its covariance

The correlation function $\xi(r_{\perp}, r_{\parallel})$ was measured for the three continuum methods. Figure 5 shows the result using the C2 method, averaged for three ranges of $\mu = r_{\parallel} / \sqrt{r_{\parallel}^2 + r_{\perp}^2}$. (The

analogous plots for C1 and C3 are in appendix C.) The superimposed curves present the results of fits as described in section 5. The full curve displays the best fit while the dashed curves are the fit when the parameters α_{\perp} and α_{\parallel} (equation 11) are set to unity, i.e. imposing the BAO peak position of the fiducial cosmology.

We have evaluated the covariance matrix, $C(r_{\perp}, r_{\parallel}, r'_{\perp}, r'_{\parallel})$ using two methods described in Appendix A. The first uses subsamples and the second a Wick expansion of the four-point function of the δ field. The two methods give covariances whose differences lead to no significant differences in fits for cosmological parameters. We use the sub-sample covariance matrix in the standard fits.

The 2500×2500 element covariance matrix has a relatively simple structure. By far the most important elements are the diagonal elements which are, to good approximation, inversely proportional to the number of pixel pairs used in the calculation of the correlation function:

$$C(r_{\perp}, r_{\parallel}, r_{\perp}, r_{\parallel}) \sim \frac{0.041}{N_{\text{pairs}}} . \quad (8)$$

This is about twice the value that one would calculate assuming all pixel pairs used to calculate $\xi(r_{\perp}, r_{\parallel})$ are independent. This decrease in the effective number of pixels is due to the correlations between neighboring pixels on a given quasar: because of these correlations, a measurement of $\xi(r_{\perp}, r_{\parallel})$ using a pair of pixels from two quasars is not independent of another measurement of $\xi(r_{\perp}, r_{\parallel})$ using the same two quasars.

The off-diagonal elements of the covariance matrix also have a simple structure. The reasons for this structure is made clear by the Wick expansion in Appendix A, which relates the covariance to correlations within pairs-of-pairs of pixels. The strongest correlations are in pairs-of-pairs where both pairs involve the *same* two quasars. To the extent that two neighboring forests are parallel, these terms contribute only to the covariance matrix elements with $r_{\perp} = r'_{\perp}$, corresponding to the transverse separation of the forests. The elements of the correlation matrix as a function of $r_{\parallel} - r'_{\parallel}$ are illustrated in figure 6 (top left); they follow closely the correlation function $\xi(\Delta\lambda)$ found within individual forests.

The covariance for $r_{\perp} \neq r'_{\perp}$ is due to pairs-of-pairs involving three or more quasars and, for small $|r_{\perp} - r'_{\perp}|$, to the fact that neighboring forests are not exactly parallel. As illustrated in figure 6, the covariances are rapidly decreasing functions of $r_{\parallel} - r'_{\parallel}$ and $r_{\perp} - r'_{\perp}$.

The statistical precision of the sub-sampling method is ~ 0.02 for individual elements of the correlation matrix. We adopt this method for the standard analysis because it is much faster than the more precise Wick method and is therefore better adapted to studies where the data sample and/or analysis protocol is varied. Figure 6 shows that only correlations with $\Delta r_{\perp} = 0$, $\Delta r_{\parallel} < 20 h^{-1}\text{Mpc}$ are greater than the statistical precision and therefore sufficiently large for individual matrix elements to be measured accurately by sub-sampling. We therefore use the average correlations as a function of $r_{\perp} - r'_{\perp}$ and $r_{\parallel} - r'_{\parallel}$, ignoring small observed variations with r'_{\perp} and r'_{\parallel} . The analysis of the mock spectra (section 5.2) indicates that this procedure is sufficiently accurate to produce reasonable χ^2 values and that the distribution of estimated BAO peak positions is near that expected from the uncertainties derived from the χ^2 surfaces.

5. Fits for the peak position

To determine the position of the BAO peak in the transverse and radial directions, we fit the measured $\xi(r_{\perp}, r_{\parallel})$ using the techniques described in Kirkby et al. (2013).

5.1. BAO model

We fit the measured $\xi(r_{\parallel}, r_{\perp})$ to a form that includes a cosmological correlation function ξ_{cosmo} and a ‘‘broadband’’ function ξ_{bb} that takes into account imperfect knowledge of the non-BAO cosmology and distortions introduced by the analysis:

$$\xi(r_{\parallel}, r_{\perp}, \alpha_{\parallel}, \alpha_{\perp}) = \xi_{\text{cosmo}}(r_{\parallel}, r_{\perp}, \alpha_{\parallel}, \alpha_{\perp}) + \xi_{\text{bb}}(r_{\parallel}, r_{\perp}) . \quad (9)$$

The function ξ_{cosmo} is described as a sum of a non-BAO ‘‘smooth’’ function and a BAO ‘‘peak’’ function

$$\begin{aligned} \xi_{\text{cosmo}}(r_{\parallel}, r_{\perp}, \alpha_{\parallel}, \alpha_{\perp}) &= \\ &= \xi_{\text{smooth}}(r_{\parallel}, r_{\perp}) + a_{\text{peak}} \cdot \xi_{\text{peak}}(\alpha_{\parallel} r_{\parallel}, \alpha_{\perp} r_{\perp}) , \end{aligned} \quad (10)$$

where a_{peak} controls the amplitude of the BAO peak relative to the smooth contribution. The radial and transverse dilation factors describing the observed peak position relative to the fiducial peak position are

$$\alpha_{\parallel} = \frac{[D_H(\bar{z})/r_d]}{[D_H(\bar{z})/r_d]_{\text{fid}}} \quad \text{and} \quad \alpha_{\perp} = \frac{[D_A(\bar{z})/r_d]}{[D_A(\bar{z})/r_d]_{\text{fid}}} , \quad (11)$$

where r_d is the sound horizon at the drag epoch (defined to sufficient accuracy for each cosmology by equation (55) of Anderson et al. (2013)).

The function ξ_{cosmo} is calculated from the power spectrum using the following procedure. We model the Ly α forest power spectrum including redshift-space distortions and nonlinear effects as

$$\begin{aligned} P(k, \mu_k) &= b^2 (1 + \beta \mu_k^2)^2 \\ &\times \left[P_{\text{peak}}(k) \exp(-k^2 \Sigma^2(\mu_k)/2) + P_{\text{smooth}}(k) \right] , \end{aligned} \quad (12)$$

where $\mu_k \equiv \hat{z} \cdot \hat{k}$, b is the Ly α forest bias parameter and β is the redshift-space distortion parameter. Here, we have defined $P_{\text{peak}}(k) = P_{\text{lin}}(k) - P_{\text{smooth}}(k)$, where P_{lin} is the linear-theory matter power spectrum from CAMB (Lewis et al., 2000) calculated using the cosmological parameters from the first column of Table 1 and P_{smooth} is the CAMB power spectrum with the BAO feature erased following the method of Kirkby et al. (2013). The exponential function in equation (12) models the anisotropic nonlinear broadening from structure growth (Eisenstein et al., 2007) with $\Sigma^2(\mu_k) = \mu_k^2 \Sigma_{\parallel}^2 + (1 - \mu_k^2) \Sigma_{\perp}^2$ and is only applied to the BAO feature. The default values we have adopted are $\Sigma_{\parallel} = 6.41 h^{-1}\text{Mpc}$ and $\Sigma_{\perp} = 3.26 h^{-1}\text{Mpc}$, which are inferred from the amplitude of the variation of linear peculiar velocities along the line of sight that cause a relative displacement of pixel pairs contributing to the BAO peak form (White, 2014).

The power spectrum multipoles are given by

$$P_{\ell}(k) = \frac{2\ell + 1}{2} \int_{-1}^{+1} P(k, \mu_k) L_{\ell}(\mu_k) d\mu_k , \quad (13)$$

where L_{ℓ} is the Legendre polynomial. The corresponding correlation function multipoles are then

$$\xi_{\ell, \text{cosmo}}(r) = \frac{i^{\ell}}{2\pi^2} \int_0^{\infty} k^2 j_{\ell}(kr) P_{\ell}(k) dk , \quad (14)$$




Fig. 6. The correlation $C(r_{\perp}, r_{\parallel}, r'_{\perp}, r'_{\parallel})/[Var(r_{\perp}, r_{\parallel})Var(r'_{\perp}, r'_{\parallel})]^{1/2}$ as a function of $r_{\parallel} - r'_{\parallel}$ (averaged over $(r'_{\perp}, r'_{\parallel})$). The top figures are for $r_{\perp} - r'_{\perp} = 0$ over the full range of Δr_{\parallel} (left) and for $\Delta r_{\parallel} > 20 h^{-1} \text{Mpc}$ (right). The bottom two figures are for $r_{\perp} - r'_{\perp} = 4 h^{-1} \text{Mpc}$ (left) and for $r_{\perp} - r'_{\perp} = 8 h^{-1} \text{Mpc}$ (right). Shown are the correlations determined by sub-sampling and by a Wick expansion. The latter correlations are decomposed into the pair-of-pair types, T1-T6, as explained in appendix A.

where j_{ℓ} is the spherical Bessel function. Finally, the correlation function is the sum of the multipoles

$$\xi_{\text{cosmo}}(r, \mu) = \sum_{\ell=0,2,4} L_{\ell}(\mu) \xi_{\ell, \text{cosmo}}(r). \quad (15)$$

The nonlinear broadening in principle transfers power to higher even multipoles $\ell = 6, 8, \dots$, but the contribution from these higher-order multipoles is negligible.

We wish to ensure the insensitivity of our results to non-BAO cosmology and to inaccurately modeled astrophysical effects like UV fluctuations, nonlinear effects and DLAs. We therefore use a ‘‘broadband’’ function, ξ_{bb} to include inaccuracies in the non-BAO correlation function as well as distortions due, for example, to continuum fitting. We use the form

$$\xi_{\text{bb}}(r_{\parallel}, r_{\perp}) = \sum_{j=0}^{j_{\text{max}}} \sum_{i=0}^{i_{\text{max}}} a_{i,j} L_{2j}(\mu) / r^i, \quad (16)$$

where the L_{2j} is the Legendre polynomial of order $2j$. Our standard model uses $(i_{\text{max}}, j_{\text{max}}) = (2, 2)$.

The standard fits use the fiducial values of Σ_{\perp} and Σ_{\parallel} , and set $a_{\text{peak}} = 1$. They thus have four physical free parameters

$(b, \beta, \alpha_{\perp}, \alpha_{\parallel})$ and, for the fiducial model, nine broadband distortion parameters. The standard fit uses the range $40 h^{-1} \text{Mpc} < r < 180 h^{-1} \text{Mpc}$ giving a total of 1515 bins in $(r_{\parallel}, r_{\perp})$ for the correlation function measurements that are actually used in the fit, and 1502 degrees of freedom.

5.2. Fits with the mock data sets

Figure 7 summarizes the results of the cosmological fits on the 100 sets of mock spectra. The mean recovered α_{\parallel} and α_{\perp} are consistent with unity, indicating no bias in the measurement of the BAO peak position.³ The numbers of α_{\parallel} measurements within 1σ and 2σ of unity are 61 and 93, consistent with the expected numbers, 68 and 95.5. For α_{\perp} the numbers are 68 and 95. For the combined $(\alpha_{\parallel}, \alpha_{\perp})$ measurements, 70 are within the 1σ and 93 within the 2σ contours.

³ After the analysis of the mocks, it was realized that they had been analyzed with a model (the fiducial model of Table 1) that was slightly different from the model used to produce them, which had the same $\Omega_M h^2$ but $\Omega_b h^2 = 0$ instead of $\Omega_b h^2 = 0.0006$. Its value of r_d is 0.15 Mpc less than the fiducial r_d so the expected mean values of α_{\perp} and α_{\parallel} are 0.999, which is sufficiently close to unity for the precision of this study.




Fig. 7. Summary of the results of fits for $(\alpha_{\parallel}, \alpha_{\perp})$ for the 100 mock catalogs. The histograms show the best-fit values, the minimum χ^2 values and the 1σ uncertainties.




Fig. 8. Measured α_{\parallel} and α_{\perp} for the 100 mock catalogs.

The r.m.s. deviations of α_{\parallel} and α_{\perp} are 0.029 and 0.057, respectively. The mean χ^2 is near the number of degrees of freedom, indicating that the model represents sufficiently well the mock observations and that the covariance matrix is well estimated.




Fig. 9. Difference in best-fit α_{\parallel} and α_{\perp} values between high redshift ($z > 2.295$) and low redshift ($z < 2.295$) subsets of the 100 mock realizations and the observations (red star). Compared to figure 8, the plot shows the degraded precision resulting from the division of the data into two redshift bins.




Fig. 10. Constraints on $(\alpha_{\parallel}, \alpha_{\perp})$ using the three continuum estimators, C1 (red), C2 (blue) and C3 (green). The solid and dashed contours correspond to 1σ and 2σ ($\Delta\chi^2 = 2.3, 6.2$).

Figure 8 shows an anti-correlation between the recovered α_{\parallel} and α_{\perp} , with a correlation coefficient of -0.6 . The quantity of the form $\alpha_{\parallel}^w \alpha_{\perp}^{1-w}$ with the smallest mock-to-mock variance has $w \sim 0.7$, with an r.m.s. deviation of 0.017. This result is to be contrasted with the optimal quantity for galaxy surveys, $\sim \alpha_{\parallel}^{1/3} \alpha_{\perp}^{2/3}$. The difference is due to the fact that redshift distortions are more important for the Ly α forest, a consequence of the low bias factor which enables more precise measurements in the r_{\parallel} direction even though there are two dimensions for r_{\perp} and only one for r_{\parallel} . This effect is evident in Fig. 5 where the BAO peak is most easily seen for $\mu > 0.8$.

Figure 9 presents the results of fits separating the mock data into two redshift bins, $z < 2.295$ and $z > 2.295$. The differences between the measured α_{\parallel} and α_{\perp} for the two bins are typically of order 10%.

Table 2. Results for the standard fit and modified fits. The standard fit uses the C2 continuum, a broadband defined by $(i_{\max}, j_{\max}) = (2, 2)$, a forest defined by $104.0 < \lambda_{\text{rf}} < 120.0$ nm, and $a_{\text{peak}} = 1$.

analysis	α_{\parallel}	α_{\perp}	β	$b(1 + \beta)$	χ^2_{\min}/DOF
standard(C2)	$1.054^{+0.032}_{-0.031}(1\sigma)^{+0.069}_{-0.063}(2\sigma)$	$0.973^{+0.056}_{-0.051}(1\sigma)^{+0.199}_{-0.103}(2\sigma)$	1.47 ± 0.87	-0.39 ± 0.05	1499.0/(1515-13)
C1	$1.038^{+0.035}_{-0.037}(1\sigma)^{+0.073}_{-0.074}(2\sigma)$	$1.054^{+0.132}_{-0.093}(1\sigma)^{+0.246}_{-0.176}(2\sigma)$	3.47 ± 2.78	-0.42 ± 0.05	1571.5/(1515-13)
C3	$1.038^{+0.026}_{-0.039}(1\sigma)^{+0.054}_{-0.071}(2\sigma)$	$1.041^{+0.259}_{-0.063}(1\sigma)^{+0.259}_{-0.126}(2\sigma)$	2.28 ± 1.24	-0.46 ± 0.05	1603.7/(1515-13)
β -prior (1.4 ± 0.4)	$1.055^{+0.032}_{-0.031}(1\sigma)^{+0.068}_{-0.063}(2\sigma)$	$0.972^{+0.053}_{-0.051}(1\sigma)^{+0.117}_{-0.102}(2\sigma)$	1.41 ± 0.36	-0.39 ± 0.04	1499.1/(1515-13)
a_{peak} free	$1.054^{+0.035}_{-0.031}(1\sigma)^{+0.078}_{-0.063}(2\sigma)$	$0.973^{+0.057}_{-0.052}(1\sigma)^{+0.232}_{-0.104}(2\sigma)$	1.49 ± 1.08	-0.38 ± 0.24	1499.0/(1515-14)
$\Sigma_{\perp} = \Sigma_{\parallel} = 0$	$1.053^{+0.029}_{-0.028}(1\sigma)^{+0.062}_{-0.059}(2\sigma)$	$0.961^{+0.055}_{-0.052}(1\sigma)^{+0.254}_{-0.103}(2\sigma)$	1.31 ± 0.87	-0.34 ± 0.05	1501.2/(1515-13)
$\Sigma_{\perp}, \Sigma_{\parallel}$ free	$1.063^{+0.041}_{-0.036}(1\sigma)^{+0.101}_{-0.073}(2\sigma)$	$0.976^{+0.053}_{-0.05}(1\sigma)^{+0.124}_{-0.102}(2\sigma)$	1.56 ± 0.80	-0.46 ± 0.07	1497.2/(1515-15)
no special DLA treatment	$1.049^{+0.038}_{-0.034}(1\sigma)^{+0.089}_{-0.068}(2\sigma)$	$0.954^{+0.053}_{-0.049}(1\sigma)^{+0.132}_{-0.096}(2\sigma)$	0.36 ± 0.46	-0.33 ± 0.06	1489.7/(1515-13)
$104.5 < \lambda_{\text{rf}} < 118.0$ nm	$1.052^{+0.041}_{-0.041}(1\sigma)^{+0.145}_{-0.094}(2\sigma)$	unconstrained	2.37 ± 2.81	-0.34 ± 0.07	1448.2/(1515-13)
No spectra with DLAs	$1.031^{+0.035}_{-0.035}(1\sigma)^{+0.074}_{-0.074}(2\sigma)$	$1.073^{+0.117}_{-0.082}(1\sigma)^{+0.228}_{-0.171}(2\sigma)$	2.38 ± 1.91	-0.43 ± 0.06	1506.5/(1515-13)
$z < 2.295$	$0.996^{+0.052}_{-0.054}(1\sigma)^{+0.113}_{-0.134}(2\sigma)$	$0.89^{+0.064}_{-0.053}(1\sigma)^{+0.148}_{-0.108}(2\sigma)$	1.10 ± 0.92	-0.31 ± 0.06	1523.0/(1515-13)
$z > 2.295$	$1.096^{+0.037}_{-0.036}(1\sigma)^{+0.079}_{-0.073}(2\sigma)$	$0.994^{+0.057}_{-0.049}(1\sigma)^{+0.155}_{-0.1}(2\sigma)$	1.61 ± 1.05	-0.48 ± 0.06	1479.1/(1515-13)
$a_{\text{peak}} = 0$	-	-	-	-	1526.2/(1515-11)

5.3. Fits with the observations

Table 2 gives the results of fits of the data for a variety of data sets and analysis assumptions. The first line lists our standard analysis using the C2 continua:

$$\alpha_{\parallel} = 1.054^{+0.032}_{-0.031} \quad \text{and} \quad \alpha_{\perp} = 0.973^{+0.056}_{-0.051}. \quad (17)$$

The precisions on α_{\parallel} and α_{\perp} inferred from our χ^2 fitting procedure are typical of those found using the 100 mock catalogs (figure 7). The full contours presented in figure 10 show that these errors are somewhat non-Gaussian, with an anti-correlation between α_{\parallel} and α_{\perp} . In particular, the 2σ contour extend asymmetrically to large α_{\perp} , consistent with the visual impression from Figure 5. The most precisely determined combination is

$$\alpha_{\parallel}^{0.7} \alpha_{\perp}^{0.3} = 1.025 \pm 0.021. \quad (18)$$

The next seven lines of Table 2 present the results of analyzes using the standard data set but with modified assumptions: use of the non-standard continua C1 and C3; adding a Gaussian prior to the redshift distortion parameter around its nominal value $\beta = 1.4$ of width 0.4; freeing the peak amplitude a_{peak} ; fitting the nonlinearity parameters, Σ_{\parallel} and Σ_{\perp} , or setting them to zero (and thus not correcting for nonlinearities); using spectra with one or more DLAs but not giving them a special treatment (fit with Voigt profile). Because these seven fits all use the same data set, any variation of the results at the 1σ level might indicate a systematic effect. In fact, all configurations produce results that are consistent at the sub- σ level. We note however that the 1σ precision on α_{\perp} is degraded through use of C1 and C3, although C1 does almost as well as C2 at the 2σ level. The higher sensitivity of the α_{\perp} uncertainty to the continuum method compared to that for the α_{\parallel} uncertainty might be expected from the low statistical significance of the BAO peak in transverse directions. The central value and error of α_{\parallel} are robust to these differences.

The next two lines in table 2 are the results for C2 with reduced data sets: using a short forest ($104.5 < \lambda_{\text{rf}} < 118.0$ nm) further from the Ly α and Ly β peaks or removing spectra with one or more DLAs. Both results are consistent at 1σ with those obtained with the more aggressive standard data set but, as expected, with larger statistical errors.




Fig. 11. The effect of metals on the measured correlation function for 10 mock sets. The red circles show $r^2 \xi(r)$ for $\mu > 0.8$ averaged over the 10 mock sets. The blue circles show the difference between $r^2 \xi(r)$ and $r^2 \xi(r)$ in the same mock realization but without metals. The light red and blue lines show the results for individual mock sets and the error bars give the standard deviation of the 10 realizations.

The next two lines present the results obtained by dividing the pixel-pair sample into two redshift bins. The two results now correspond to fairly independent samples and are in agreement at the 2σ level. The differences between measurements of $(\alpha_{\parallel}, \alpha_{\perp})$ for the two redshift bins in the mock spectra are displayed in figure 9, suggesting that the observed difference is not atypical of that observed with the mock spectra.

The last line of table 2 is the χ^2 for a fit with no BAO peak. Comparison with the first line reveals $\Delta\chi^2 = 27.2$ for two extra degrees of freedom, corresponding to a 5σ detection.

6. Systematic Errors

The uncertainties reported in Table 2 are statistical and are derived from the χ^2 surface. In this section we discuss possible




Fig. 12. Values α_{\parallel} (blue dots) and α_{\perp} (red dots) recovered from the DR11 data for different choices of the minimum transverse separation, r_{\perp}^{\min} , used in the fit. The measured values do not change significantly when eliminating the small r_{\perp} bins that may be contaminated by correlations due to absorption by metals.

Table 3. Fit results with the C2 continuum with a modified fitting range [standard: $40 < r < 180 h^{-1}\text{Mpc}$] and number of terms in the broadband (equation 16) [standard: $(i_{\max}, j_{\max}) = (2, 2)$].

	α_{\parallel}	α_{\perp}	χ_{\min}^2/DOF
standard (C2)	1.054 ± 0.032	0.973 ± 0.055	1499.0/(1515-13)
range ($h^{-1}\text{Mpc}$)			
$60 < r < 180$	1.045 ± 0.032	0.986 ± 0.063	1391.8/(1415-13)
$40 < r < 160$	1.052 ± 0.033	0.974 ± 0.053	1139.2/(1177-13)
(i_{\max}, j_{\max})			
(2,3)	1.057 ± 0.032	0.970 ± 0.050	1484.2/(1515-16)
(3,2)	1.050 ± 0.033	0.987 ± 0.067	1497.8/(1515-16)
(3,3)	1.051 ± 0.034	0.986 ± 0.068	1479.2/(1515-20)

systematic errors. We find no evidence for effects that add uncertainties comparable to the statistical errors.

In this paper, we derive cosmological information by comparing the measured flux correlation function with a model, defined by equations (9), (10) and (16), that depends on cosmological parameters (equation 10). Systematic errors in the derived parameters can result if either the assumed model or the measured correlation function differ systematically from the true flux-correlation function.

In fitting the data, we add to the assumed cosmological correlation function ξ_{cosmo} a general “broadband” form ξ_{bb} (equation 16). The role of ξ_{bb} is to make the fit sensitive only to the position of the BAO peak and not to the more uncertain smooth component of ξ_{cosmo} . To verify that the broadband does indeed remove any sensitivity to smooth components of the correlation function, we varied the form of the assumed broadband and the range over which it is fit. The results, listed in table 3, show no significant variation of the derived $(\alpha_{\parallel}, \alpha_{\perp})$, indicating that the broadband is performing as required. Of particular significance, adding greater freedom to ξ_{bb} has only a $\sim 10\%$ impact on the size of the α_{\parallel} error, although it has a larger impact (20 – 30%) on the α_{\perp} error.

Because the use of ξ_{bb} makes our results insensitive to smooth features in ξ , we are primarily concerned with *non-smooth* effects either due to observing or analysis artifacts or to physical effects invalidating the assumed theoretical form (equation 9).

We first consider errors in the theoretical form of the correlation function. The function ξ_{cosmo} in (9) is subject to uncertainties arising from nonlinear effects and, more importantly, in the astrophysical processes that determine the flux transmission correlations from matter correlations. The resulting uncertainties in the dominant Ly α absorption would be expected to generate only errors that vary slowly with r and are therefore absorbed into ξ_{bb} . On the other hand, absorption by metals, not included in (9), generates excess correlation in individual forests in the form of narrow peaks centered at the wavelength separations between $\lambda_{\text{Ly}\alpha}$ and metal lines. For example, the SiII(1260.42) absorption correlated with Ly α (1215.67) absorption gives rise to a narrow peak of excess correlation at $r = 110 h^{-1}\text{Mpc}$ on the line of sight, at $z = 2.34$. This narrow peak is smeared because of the range of observed redshifts, over a width $\Delta r \sim \pm 5 h^{-1}\text{Mpc}$. This correlation in the absorption in individual quasar spectra induces a correlation in the spectra of neighboring quasars (small r_{\perp}), because they probe correlated structures of Ly α and SiII absorption in the IGM.

As described in Bautista et al. (in preparation), we added metals to the mock spectra to estimate their importance. As expected, the mocks indicate that the metal-induced correlation falls off rapidly with transverse separation, dropping by a factor of five between the first and third r_{\perp} bin. Figure 11 shows the effect on ten sets of mocks in the important region $\mu > 0.8$. The modifications do show structure that might not be modeled by our broadband term. Fortunately, the effect has no significant impact on the measured position of the BAO peak, with $(\alpha_{\parallel}, \alpha_{\perp})$ for metaled and metal-free mocks having a mean difference and mock-to-mock dispersion of $\Delta\alpha_{\parallel} = 0.002 \pm 0.003$ and $\Delta\alpha_{\perp} = 0.003 \pm 0.009$.

Because the amplitude of the metal lines is somewhat uncertain we have empirically verified that they are unimportant by re-performing the fit of the DR11 data after excising the correlation function bins with $r_{\perp} < r_{\perp}^{\min}$. Most of the metal correlation occurs within $r_{\perp} < 10 h^{-1}\text{Mpc}$, so the presence of any unexpected effect from metal lines would have to be made apparent by a dependence of the BAO results on r_{\perp}^{\min} . In fact, the results are remarkably stable, as shown in Figure 12. We thus conclude that absorption by metals is unlikely to significantly affect the measured position of the BAO peak.

We now turn to artifacts introduced by the analysis. The measured correlation function will be different from the true flux-transmission correlation function because of systematic errors in the flux-transmission field, $\delta_q(\lambda)$, defined by equation (2). Such errors can be introduced through an inaccurate flux-calibration or an inaccurate estimate of the function $C_q(\lambda)\bar{F}(z)$. These errors in $\delta_q(\lambda)$ will generate systematic errors in the correlation function if the neighboring quasars have *correlated* systematic errors.

The most obvious error in the $\delta_q(\lambda)$ arises from the necessity of using a spectral template to estimate the continuum. The C1 and C2 methods use a unique template that is multiplied by a linear function to fit the observed spectrum. This approach results in two systematic errors on the correlation function. First, as previously noted, fitting a linear function for the continuum effectively removes broadband power in individual spectra. This error will be absorbed into ξ_{bb} and not generate biases in the BAO peak position. Second, $\delta_q(\lambda)$ along individual lines of sight

will be incorrect because non-smooth quasar spectral diversity is not taken into account by the universal template. However, because the peculiarities of individual quasar spectra are determined by local effects, they would not be expected to be correlated between neighboring quasars. The tests with the mock catalogs which include uncorrelated spectral diversity confirm that the imprecisions of the continuum estimates do not introduce biases into the estimates of the BAO peak positions.

Errors introduced by the flux calibration are potentially more dangerous. The BOSS spectrograph (Smee et al., 2012) is calibrated by observing stars whose spectral shape is known. Most of these objects are F-stars whose spectra contain the Balmer series of hydrogen lines. The present BOSS pipeline procedure for calibration imperfectly treats the standard spectra in the neighborhood of the Balmer lines, resulting in calibration vectors, $C(\lambda)$, that show peaks at the Balmer lines of amplitude $\langle \Delta C/C \rangle \sim 0.02 \pm 0.004$, where the ± 0.004 refers to our estimated quasar-to-quasars r.m.s. variation of the Balmer artifacts (Busca et al., 2013). If uncorrected, these calibration errors would lead to $\delta \sim 0.02$ at absorber redshifts corresponding to the Balmer lines. The subtraction of the mean δ -field, $\langle \delta(\lambda) \rangle$ in our analysis procedure removes this effect on average but does not correct calibration vectors individually. Because of the relative uniformity of the Balmer feature in the calibration vectors, this mean correction is expected to be sufficient. In particular, we have verified that no significant changes in the correlation function appear when it is calculated taking into account the observed correlations in the Balmer artifacts, $\Delta C/C$.

To verify this conclusion, we have searched for Balmer artifacts in the measured $\xi(r_{\parallel}, r_{\perp}, \langle \lambda \rangle)$ where $\langle \lambda \rangle$ is the mean wavelength of the pixel pair. If our mean correction is insufficient, there would be excess correlations at $r_{\parallel} = 0$ and $\langle \lambda \rangle$ equal to a Balmer wavelength. Artifacts would also appear at r_{\parallel} and $\langle \lambda \rangle$ corresponding to pairs of Balmer lines. For example, the pair [H δ (410 nm), H ϵ (397 nm)] would produce excess correlation at the corresponding radial separation $98 h^{-1} \text{Mpc}$ and $\langle \lambda \rangle = 403 \text{ nm}$. A search has yielded no significant correlation excesses. Additionally, removing from the analysis pixel pairs near (397,410)nm, dangerously near the BAO peak, does not generate any measureable change in the BAO peak position.

7. Cosmological implications

The standard fit values for $(\alpha_{\parallel}, \alpha_{\perp})$ from Table 2 combined with the fiducial values from Table 1 yield the following results

$$\frac{D_H(2.34)}{r_d} = 9.18 \pm 0.28(1\sigma) \pm 0.6(2\sigma) \quad (19)$$

and

$$\frac{D_A(2.34)}{r_d} = 11.28 \pm 0.65(1\sigma) \pm 2.8(2\sigma). \quad (20)$$

The blue shading in Figure 13 shows 68.3% and 95.5% likelihood contours for these parameters, showing a mild anti-correlation between them. These constraints can be expressed equivalently as

$$H(z = 2.34) = (222 \pm 7 \text{ km s}^{-1} \text{ Mpc}^{-1}) \times \frac{147.4 \text{ Mpc}}{r_d}$$

$$D_A(z = 2.34) = (1662 \pm 96 \text{ Mpc}) \times \frac{r_d}{147.4 \text{ Mpc}}, \quad (21)$$

where we have scaled by the value $r_d = 147.4 \text{ Mpc}$ from the Planck+WP model in Table 1.




Fig. 13. Constraints on $(D_A/r_d, D_H/r_d)$. Contours show 68.3% ($\Delta\chi^2 = 2.3$) and 95.5% ($\Delta\chi^2 = 6.2$) contours from the Ly α forest auto-correlation (this work, blue), the quasar Ly α forest cross-correlation (Font-Ribera et al., 2013) (red), and the combined constraints (black). The green contours are CMB constraints calculated using the Planck+WP+SPT+ACT chains (Ade et al., 2013) assuming a flat Λ CDM cosmology.




Fig. 14. Constraints on the Ω_{Λ} CDM parameters $(\Omega_{\Lambda}, \Omega_M)$ based on the auto-correlation contours of figure 13. The contours show 68.3% and 95.5% confidence levels. The Planck value of $\Omega_B h^2$ is assumed together with a Gaussian prior for $H_0 = 70.6 \pm 3.2 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The yellow star is the Planck Λ CDM measurement and the dashed line corresponds to a flat universe.

Our measured values of D_H/r_d and D_A/r_d (equations 19 and 20) can be compared with those predicted by the two CMB inspired flat Λ CDM models from Table 1: (8.570, 11.76) for Planck+WP and (8.648, 11.47) for WMAP9+ACT+SPT. Figure 13 demonstrates that our values differ by 1.8σ from those of the Planck+WP model. They differ from the WMAP9+ACT+SPT model by 1.6σ . We emphasize that, in contrast to the values of α_{\parallel} and α_{\perp} , the constraints quoted in equations (19)-(21) are independent of the fiducial model adopted in the analysis, at least over a substantial parameter range. We have confirmed this expectation by repeating some of our analyses using the Planck+WP parameters of Table 1 in place of our standard fiducial model, finding negligible change in the inferred values of D_H/r_d and D_A/r_d .

To illustrate this tension, we show in figure 14 values of Ω_M and Ω_{Λ} that are consistent with our measurements. We consider

models of Λ CDM with curvature, having four free parameters: the cosmological constant, the matter and the baryon densities, and the Hubble constant ($\Omega_\Lambda, \Omega_M, \Omega_B h^2, h$). For clarity, we place priors on two of them, adopting the Planck value of $\Omega_B h^2$ (Ade et al., 2013) and adopting a wide prior on $h = 0.706 \pm 0.032$ meant to cover the value measured with the local distance ladder (Riess et al., 2011) and that measured with CMB anisotropies assuming a Λ CDM cosmology (Ade et al., 2013). Figure 14 shows the resulting constraints. We see that for flat Λ CDM models, it is necessary to lower Ω_M from ~ 0.3 to ~ 0.2 to produce our measured D_H/r_c and D_A/r_d .

The tension with CMB data is also seen in the BAO measurement using the quasar-Ly α forest cross-correlation (Font-Ribera et al., 2013). Red contours in Figure 13 show their 68.3% and 95.5% likelihood contours. The implied values of D_A and D_H are consistent between the auto-correlation and cross-correlation measurements, but the statistical errors are interestingly complementary. The auto-correlation constrains D_H more tightly than D_A because redshift-space distortions are so strong in Ly α forest clustering, a consequence of the low bias factor of the forest. While there are far fewer quasar-forest pairs than forest-forest pairs, the cross-correlation still yields a useful BAO signal because the quasars themselves are highly biased, which boosts the clustering amplitude. However, redshift-space distortions are weaker in the cross-correlation for the same reason, and the cross-correlation analysis yields comparable statistical errors in the transverse and line-of-sight BAO. The cross-correlation constraint therefore suppresses the elongated tails of the auto-correlation likelihood contours seen in Figure 10 towards high D_A and correspondingly low D_H .

The statistical errors in these BAO measurements are dominated by combinations of limited sampling of the volume probed, by instrumental noise in the Ly α forest spectra, and (for the cross-correlation measurement) by shot noise of the quasar density field. For this reason, the statistical errors in the two BAO measurements are almost completely uncorrelated, as discussed further in Appendix B. We have therefore combined the two likelihood surfaces as independent to produce the joint likelihood contours shown by the solid lines in Figure 13. Marginalized 1-d constraints from the combined likelihood are:

$$\frac{D_H(2.34)}{r_d} = 9.15^{+0.20}_{-0.21} (1\sigma) \quad {}^{+0.40}_{-0.42} (2\sigma) \quad (22)$$

and

$$\frac{D_A(2.34)}{r_d} = 10.93^{+0.35}_{-0.34} (1\sigma) \quad {}^{+0.75}_{-0.65} (2\sigma). \quad (23)$$

These numbers can be compared with the green contours in Figure 13, which show the 68.3%, 95.5%, and 99.7% confidence contours on D_A/r_d and D_H/r_d derived from CMB data (specifically, using the Planck + WMAP polarization + SPT + ACT chains available from the Planck Collaboration; this data set gives results very similar to the Planck+WP model of Table 1), assuming a flat Λ CDM cosmological model. These predictions lie outside the 95.5% likelihood interval for the combined cross- and auto-correlation BAO measurements, an $\approx 2.5\sigma$ tension with the data. In more detail, the Λ CDM prediction is approximately 2σ below the value of D_H inferred from the auto-correlation and approximately 2σ above the value of D_A inferred from the cross-correlation, deviations of $\approx 7\%$ in each case. The tension between the CMB-constrained flat Λ CDM model and the auto-correlation measurement of D_H is evident in the top panel of Figure 5, where the peak in the data is visually to the left of

the peak in the fiducial model (and would be even more to the left of the Planck+WP model).

How seriously should one take this tension? Concerning the auto-correlation function, the success of our method in reproducing the correct parameters when averaged over our 100 mock catalogs, and the insensitivity of our derived α_{\parallel} and α_{\perp} to many variations of our analysis procedure as discussed in §6, both suggest that systematic biases in our measurements should be less than our quoted errors. The agreement between the directly-estimated statistical errors and the scatter in best-fit α values for our mock catalogs indicate that our error estimates themselves are accurate, although with 100 mock catalogs we cannot test this accuracy stringently. The most significant impact seen with varying the analysis procedure in §6 is the larger statistical errors on α_{\parallel} and α_{\perp} for the continuum subtraction method C1. Detailed examination of the likelihood contours in Figure 10 shows that the larger α_{\parallel} errors (and lower central value) for the C1 method are a consequence of its weaker constraint on α_{\perp} , which allows contours to stretch into the region of low α_{\parallel} and high α_{\perp} . These regions are inconsistent with the stronger α_{\perp} constraints of the cross-correlation measurement, so in a joint likelihood they would be eliminated in any case. Furthermore, our standard C2 subtraction method is clearly more realistic than the C1 method because it is based on a more realistic flux PDF rather than a Gaussian approximation to it. Nonetheless, the variation seen in Table 2 suggests some degree of caution about the precision of our statistical errors, even though we have no clear evidence that they are underestimated, particularly because of the relatively weak detection of the transverse BAO signal.

While it is premature to conclude that a major modification of Λ CDM is needed, it is nevertheless interesting to note what sort of changes are indicated by the data. The most widely discussed extensions to flat Λ CDM, allowing non-zero space curvature or a dark energy equation-of-state with $w \neq -1$, do not readily resolve the tension seen in Figure 13 without running afoul of other constraints. This is because of the necessity of decreasing $D_A(2.34)$ while increasing $D_H(2.34)$, something not easily done since the former is related to the integral of the latter.

Requirements for more general forms of dark energy can be found by considering our measurement of $H(z)$, which, combined with the Friedman equation, determines the density of dark energy $\rho_{\text{de}}(z)$. Assuming space to be flat and matter to be conserved, and neglecting the radiation density, we have

$$\frac{8\pi G}{3} \rho_{\text{de}}(z) = H^2(z) - H_0^2 \Omega_M (1+z)^3. \quad (24)$$

Dividing by $\rho_{\text{de}}(z=0)$ gives

$$\frac{\rho_{\text{de}}(z)}{\rho_{\text{de}}(z=0)} = \frac{H(z)^2 - \Omega_M H_0^2 (1+z)^3}{(1 - \Omega_M) H_0^2}. \quad (25)$$

The uncertainty on $\rho_{\text{de}}(z=2.34)/\rho_{\text{de}}(0)$ is dominated by the difference between two large numbers in the numerator. If we use the precise values of r_d and $\Omega_M h^2 = 0.143 \pm 0.003$ from the Planck+WP CMB power spectrum measurement, the uncertainty is dominated by that of our value of $r_d H(z=2.34)$ (eqn. 22). We find

$$\frac{\rho_{\text{de}}(z=2.34)}{\rho_{\text{de}}(z=0)} = -1.2 \pm 0.8. \quad (26)$$

The difference of $\sim 2.5\sigma$ from the expected value of unity for the Λ CDM model is the same as the difference discussed above (although the quoted error above implies a 2.8σ deviation, this

is reduced slightly by the non-gaussianity of the likelihood distribution of the measured D_H). If a negative value of ρ_{de} were to persist as measurement errors on $H(z)$ from BAO and $\Omega_M h^2$ from the CMB are improved, this would imply that the dark energy density at $z=2.4$ is less than that of $z=0$, perhaps even with the opposite sign, that matter was not conserved from the epoch of recombination, or that the universe is closed.

8. Conclusions

The Ly α correlation data presented in this study constrain D_H/r_d and D_A/r_d at $z \sim 2.34$.⁴ The 3.0% precision on D_H/r_d and 5.8% precision on D_A/r_d obtained here improve on the precision of previous measurements: 8% on D_H/r_d (Busca et al., 2013); and 3.4% on D_H/r_d and 7.2% on D_A/r_d (Slosar et al., 2013). The increasing precision of the three studies is primarily due to their increasing statistical power, rather than to methodological improvements. The 2% precision on the optimal combination $D_H^{0.7} D_A^{0.3}/r_d$ can be compared with the 1% precision for $D_V(z = 0.57)/r_d$ obtained by Anderson et al. (2013).

The derived values of D_H/r_d and D_A/r_d obtained here with the Ly α auto-correlation are similar to those inferred from the Quasar-Ly α -forest cross-correlation (Font-Ribera et al., 2013), as shown in figure 13. At the two-standard-deviation level, the two techniques are separately compatible with the ‘‘Planck+WP’’ and fiducial models of Table 1. However, the combined constraints are inconsistent with the Planck+WP Λ CDM model at $\approx 2.5\sigma$ significance, given our estimated statistical uncertainties. The tests presented in earlier sections suggest that our statistical error estimates are accurate and that systematic uncertainties associated with our modeling and analysis procedures are smaller than these statistical errors. However, we have not shown that these systematic uncertainties are completely negligible, and further investigations with larger numbers of carefully tailored mock catalogs will be needed to better quantify them.

The Ly α -forest BAO measurements are statistically unlikely in a flat Λ CDM universe constrained by the CMB, although at a level that is still insufficient to be confidently interpreted as a real discrepancy. The cosmological implications of our results will be investigated in much greater depth in a forthcoming paper (BOSS, in preparation) where we combine the Ly α -forest BAO with the BOSS galaxy BAO results at lower redshift and with CMB and supernova data, which enables interesting constraints on a variety of theoretical models.

Acknowledgements. Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy Office of Science. The SDSS-III web site is <http://www.sdss3.org/>.

SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of

⁴ The *baofit* software used in this paper is publicly available at <http://github.com/deepzot/baofit/>. The measured cross-correlation function and its covariance matrix, and the instructions to reproduce the main BAO results presented in this paper, can be downloaded from <http://darkmatter.ps.uci.edu/baofit/>, together with the likelihood surfaces used to generate the contours in Figure 13.

Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

The French Participation Group of SDSS-III was supported by the Agence Nationale de la Recherche under contracts ANR-08-BLAN-0222 and ANR-12-BS05-0015-01.

Timothée Delubac and Jean-Paul Kneib acknowledge support from the ERC advanced grant LIDA

Matthew Pieri has received funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement n [PIIF-GA-2011-301665].

The authors acknowledge the support of France Grilles for providing computing resources on the French National Grid Infrastructure.

Appendix A: Covariance Matrix

A.1. Estimation of the covariance via a Wick expansion.

The Wick expansion for the covariance between ξ for two bins A and B is the sum over pairs of pairs:

$$C_{AB} = S_{AB}^{-1} \sum_{i,j \in A} \sum_{k,l \in B} w_i w_j w_k w_l [\xi_{ik} \xi_{jl} + \xi_{il} \xi_{jk}] \quad (\text{A.1})$$

with

$$\xi_{ij} = \langle \delta_i \delta_j \rangle. \quad (\text{A.2})$$

The pairs (i, j) and (k, l) refer to the ends of the vectors $r_A \in A$ and $r_B \in B$. The normalization factor is

$$S_{AB} = \sum_{i,j \in A} (w_i w_j) \sum_{k,l \in B} (w_k w_l). \quad (\text{A.3})$$

As illustrated in figure A.1, there are six types of pairs-of-pairs, $(ijkl)$, characterized by the number of distinct points $(2,3,4)$ and numbers of quasars $(2,3,4)$.




Fig. A.1. The six types of pairs of pairs. The dashed lines refer to quasar lines-of-sight. The variances are dominated by types 1, 2 and 3. The $(r_{\perp} - r'_{\perp} = 0)$ covariances are dominated by types 2 and 3.

The complete sum of pairs-of-pairs would require a prohibitively large amount of computer time. We have therefore evaluated the sum by using only a random sample of pairs-of-pairs and by replacing products of distinct pixels with the previously evaluated correlation function, either 1d for pairs involving only one quasar or 3d for pairs involving two quasars

The variances, equation 8, are dominated ($\sim 97\%$) by the two-quasar diagrams in figure A.1. About 60% of the variance is produced by the diagonal diagram ($i = k, j = l$). The non-diagonal terms, ($i = k, j \neq l$) and ($i \neq k, j = l$) account for respectively 25% and 15% of the variance. The dominant covariances, i.e., those with $r_{\perp} = r'_{\perp}$ and $0 < r_{\parallel} - r'_{\parallel} < 20 h^{-1} \text{Mpc}$, are dominated by the non-diagonal two-quasar diagrams.

The Wick results for the important covariance matrix elements are summarized in figure 6.

A.2. Estimation of the covariance via sub-sampling.

We used a sub-sample method to estimate the covariance matrix. The method consists in organizing the space of pairs of quasars into *sub-samples*. We took advantage of the fact that quasars are observationally tagged with the number of the plate on which they were observed. A given pair belongs to the sub-sample p if the quasar with the smaller right ascension in the pair was observed at plate p . Thus there are as many sub-samples as the number of plates (N_{plates}) that compose the data sample ($N_{\text{plates}} = 2044$ for DR11).

In terms of this partition of the data sample into sub-samples, we write our estimator of the correlation function in equation 3 as:

$$\hat{\xi}_A = \frac{1}{\sum_{p=1}^{N_{\text{plates}}} w_A^p} \sum_{p=1}^{N_{\text{plates}}} \hat{\xi}_A^p w_A^p, \quad (\text{A.4})$$

where $\hat{\xi}_A^p$ is the correlation function calculated using only pairs belonging to the sub-sample p and w_A^p is the sum of their weights. The denominator is equal to the sum of weights in A , the normalization in equation 3.

Our partitioning scheme ensures that a pair of quasars contributes to one and only one sub-sample. This approach implies that the correlation between $\hat{\xi}_A^p$ and $\hat{\xi}_A^{p'}$ (with $p \neq p'$) is given only by terms of the form T4, T5 and T6. In what follows we will neglect this small correlation.

The covariance matrix is then given by:

$$\langle \hat{\xi}_A \hat{\xi}_B \rangle - \langle \hat{\xi}_A \rangle \langle \hat{\xi}_B \rangle = S_{AB}^{-1} \sum_{p=1}^{N_{\text{plates}}} w_A^p w_B^p \left(\langle \hat{\xi}_A^p \hat{\xi}_B^p \rangle - \langle \hat{\xi}_A \rangle \langle \hat{\xi}_B \rangle \right), \quad (\text{A.5})$$

where, as anticipated, we have assumed that crossed terms from different plates are zero. The final step is to use the following estimator for the expression in parentheses, the covariance within a plate:

$$\hat{C}_{AB}^p = \hat{\xi}_A^p \hat{\xi}_B^p - \hat{\xi}_A \hat{\xi}_B. \quad (\text{A.6})$$

Appendix B: Combining the results with Font-Ribera et al. (2013)

In this appendix we discuss the level of correlation between the BAO measurement presented in this paper and that measured in Font-Ribera et al. (2013) from the cross-correlation of the Ly α forest with the quasar density field, also using the DR11 of BOSS.

If both analyses were limited by cosmic variance there would be no gain in combining them, since both would be tracing the same underlying density fluctuations. However, as shown in appendix B of Font-Ribera et al. (2013), cosmic variance is only a minor contribution to the uncertainties in both measurements.

The accuracy of the Ly α auto-correlation measurement (presented here) is limited by the ‘‘aliasing noise’’ (McDonald & Eisenstein, 2007; McQuinn & White, 2011) and the instrumental noise, while the cross-correlation measurement (Font-Ribera et al., 2013) is also limited by the ‘‘shot-noise’’ of the quasar field. Since the dominant sources of fluctuation in the two measurements have a completely different nature, the cross covariance should be small.

To better quantify this statement, we calculate the covariance between the two measurements by computing the cross-correlation coefficient between a bin measured in the auto-correlation $\hat{\xi}_A$ and a bin measured in the cross-correlation $\hat{\xi}_a$, defined as:

$$r_{Aa} = \frac{C_{Aa}}{\sqrt{C_{AA} C_{aa}}}, \quad (\text{B.1})$$

where C_{AA} is the variance in the auto-correlation bin A , C_{aa} is the variance in the cross-correlation bin a , and C_{Aa} is the covariance between the two bins.




Fig. B.1. The three types of diagrams contributing to the covariance between a bin A in the Ly α auto-correlation and a bin a in the cross-correlation with quasars. The dashed lines refer to Ly α forests and the dots to quasars. The solid lines refer to Ly α pixel pairs or quasar-Ly α pairs used to measure the auto- or cross-correlation.

We calculated the covariance C_{Aa} using a Wick expansion similar to that computed in appendix A. In this case, we must compute a four-point function with three Ly α pixels and a quasar position, and the different contributions to the covariance will be products of the Ly α auto-correlation function between two pixels and the Ly α -quasar cross-correlation between a quasar and a pixel. As shown in figure B.1, there will be three types of contribution to the covariance, arising from configurations with two, three and four quasars.

The correlation of pixels in different lines of sight will in general be smaller than the correlation of pixels in the same line of sight. Therefore, we expect the right most diagram in figure B.1 to have a small contribution, since it involves pixel pairs from different lines of sight.

Direct computation shows that the contribution from three quasar diagrams is about a factor of ten larger than that from two-quasar diagrams for $r_{\perp}^A = r_{\perp}^a$. The two quasar contribution for $r_{\perp}^A \neq r_{\perp}^a$ is zero.

In figure B.2 we show the cross-correlation coefficients computed from the three-quasar diagrams as a function of $r_{\parallel}^A - |r_{\parallel}^a|$ for $r_{\perp}^A = r_{\perp}^a$ (it decreases with increasing $|r_{\parallel}^A - r_{\parallel}^a|$). As expected, the correlation between the two measurements is small, justifying the combined contours presented in figure 13.




Fig. B.2. Cross-correlation coefficients as a function of $r_{\parallel}^A - |r_{\parallel}^a|$, computed from the three-quasar diagrams. A and a refer, respectively, to bins for the auto-correlation and cross-correlation measurements.

Appendix C: Correlation function for C1, C2 and C3

Figure C.1 shows the correlation function found using the three continuum estimators in three ranges of μ (the same as those in figure 5). All three methods show a clear BAO peak in the nearly radial bin, $\mu > 0.8$. While the peak in this bin is at the same position for all continua, the absolute value of the correlation function is quite different. This is because the $\mu > 0.8$ bin is strongly affected by the distortions induced by the continuum estimator.

References

- Aardrock, C. et al. in preparation
Ade, P. A. R., N. Aghanim, C. Armitage-Caplan, et al. 2013, arXiv:1303.5076
Ahn, C.P., R. Alexandroff, C. Allende Prieto et al. 2012, ApJS, 203, 21
Ahn, C.P., R. Alexandroff, C. Allende Prieto et al. 2013, arXiv:1307.7735
Anderson, L., E. Aubourg, S. Bailey et al. 2012, MNRAS, 427, 3435
Anderson, L., E. Aubourg, S. Bailey et al. 2012, arXiv:1303.4666
Bautista, J. et al. 2014, in preparation
Becker, R.H., R.L. White & D.J. Helfand 1995, ApJ, 450, 559B
Beutler F. et al. 2011, MNRAS, 416, 3017
Blake C. et al. 2011, MNRAS, 415, 2892
Bolton A. et al. 2012, AJ, 144, 144
Bovy, J. et al. 2011, ApJ, 729, 141
Bovy, J. et al. 2012, ApJ, 749, 41
Busca, N.G. et al. 2013, A&A, 552, 96
Calabrese, E. et al. 2013, Phys. Rev. D, 87, 103012
Chuang, Chia-Hsun & Yun Wang 2012, MNRAS, 426, 226
Cole, S. et al. 2005, MNRAS, 362, 505
Dawson, K., D. Schlegel, C. Ahn et al. 2013, AJ, 145, 10
Eisenstein D. J., Zehavi, I., Hogg, D.W. et al. 2005, ApJ, 633, 560
Eisenstein, D.J. 2007, ApJ, 664, 675
Eisenstein, D.J. et al. 2011, AJ, 142, 72
Font-Ribera, A., P. McDonald & J. Miralda Escudé 2012, J. Cosmology Astropart. Phys., 01, 001
Font-Ribera, A. & Miralda-Escud, J. 2012, J. Cosmology Astropart. Phys., 07, 028
Font-Ribera, A., D. Kirkby, N. Busca et al. 2013, arXiv:1311.1767, submitted to J. Cosmology Astropart. Phys.
Fukugita, M., Ichikawa, T., Gunn, J. E., et al. 1996, AJ, 111, 1748
Gunn, J.E., et al. 2006, AJ, 131, 2332
Hamilton, A.J.S. 1992, ApJ, 385L, 5
Eyal A. Kazin, E.A., A. G. Sanchez, A. J. Cuesta, et al. 2013, arXiv:1303.4391
Kirkby, D., D. Margala, A. Slosar et al. 2013, J. Cosmology Astropart. Phys., 03, 024
Kirkpatrick, J.A., D. J. Schlegel, N. P. Ross, et al., 2011, AJ, 743, 125

- Lawrence, A., S.J. Warren, O. Almaini et al. 2007, MNRAS, 379, 1599
Lee, K.G., N. Suzuki and D. N. Spergel 2012, AJ, 143, 51
Lee, K.G., S. Bailey, L.E. Bartsch et al. 2013, ApJ, 145, 69
Lewis, A., A. Challinor & A. Lasenby 2000, ApJ, 538, 473
Martin, D.C., J. Fanson, D. Schiminovich et al. 2005, ApJ, 619, L1
McDonald, P. 2003, ApJ, 585, 34
McDonald, P., U. Seljak, D.J. Schlegel et al. 2006, ApJS, 163, 80
McDonald, P. & D. Eisenstein 2007, Phys. Rev. D, 76, 063009
McQuinn, M. & M. White 2011, MNRAS, 415, 2257
Mehta, K.T., A. Cuesta, X. Xu et al. 2012, MNRAS, 427, 2168
Noterdaeme, P., P. Petitjean, W.C. Carithers et al. 2012, A&A, 547, L1
O'Connell, R. et al. in preparation
Padmanabhan, N. et al. 2009, Phys. Rev. D, 79, 3523
Padmanabhan, N., X. Xu, D.J. Eisenstein et al. 2012, MNRAS, 427, 2132
Palanque-Delabrouille, N., C. Yèche, A. D. Myers et al. 2011, A&A, 530, 122
Pâris, I., P. Petitjean, E. Aubourg et al. 2012, A&A, 548, 66
Pâris, I., P. Petitjean, E. Aubourg et al. 2013, arXiv:1311.4870
Percival, W.J., B.A. Reid, D.J. Eisenstein et al. 2010, MNRAS, 401, 2148
Pieri, M. M., S. Frank, D. H. Weinberg, S. Mathur, and D. G. York 2010, ApJ, 724, L73
Pieri, M. M., et al. 2013, arXiv:1309.6768
Richards, G.T. et al. 2009, ApJS, 180, 67
Riess, A.G., L. Macri, S. Casertano et al. 2011, ApJ, 730, 119
Ross, N.P. et al. 2012, ApJS199, 3
Seo, H.-J., S. Ho, M. White et al. 2012, ApJ, 761, 13
Slosar, A. et al. 2011, J. Cosmology Astropart. Phys., 09, 001
Slosar, A., V. Irsic, D. Kirkby, et al. 2013, J. Cosmology Astropart. Phys., 04, 026
Smee, S., J.E. Gunn, A. Uomoto et al. 2012, arXiv:1208.2233
Suzuki, N., D. Tytler, D. Kirkman et al. 2005, ApJ, 618, 592
Suzuki, N. 2006, ApJS, 163, 110
Veropalumbo, A. et al. 2013, arXiv:1311.5895
White, M. 2003, The Davis Meeting On Cosmic Inflation. 2003 March 22-25, Davis CA., p.18
White, M. 2014, arXiv:1401.5466.
Xu, X., Cuesta, A.J., N. Padmanabhan et al. 2012, MNRAS, 431, 2834
Yèche, C., P. Petitjean, J. Rich, et al. 2009, A&A, 523, A14
York, D. G. et al. (SDSS Collaboration) 2000, AJ120, 1579




Fig. C.1. The measured correlation functions for continuum C1 (top), C2 (middle) and C3 (bottom) averaged over three angular regions: $0 < \mu < 0.5$ (left), $0.5 < \mu < 0.8$ (middle), and $\mu > 0.8$ (right). The curves show the results of fits as described in section 5. The blue dashed curve is best fit and the full red curves if the best fit for C2.